



KEPROLYTE ORAL

COMPOSITION:

Contains per ml:

Dextrose monohydrate	300	mg
Sodium chloride	100	mg
Glycine	50	mg
Sodium dihydrogen phosphate dihydrate	22	mg
Potassium chloride	13.5	mg
Sodium citrate	5	mg

DESCRIPTION:

The dextrose monohydrate sugar and the electrolytes sodium chloride, sodium dihydrogen phosphate dihydrate, potassium chloride and sodium citrate can be used for recovery of electrolyte and acid/base imbalances. The essential amino-acid glycine is added for a quicker recovery of the dehydrated animals.

INDICATIONS:

Keprolyte Oral is indicated for prevention and treatment of dehydration caused by diarrhoea in calves, cattle, goats, poultry, sheep and pigs.

SIDE EFFECTS:

None.

DOSAGE AND ADMINISTRATION:

For oral administration via drinking water. Shake well before use.

Calves, cattle, goats, sheep, pigs : 50 ml per litre of drinking water, twice daily, during 2 - 4 days.

Poultry : 100 ml per 75 - 100 litres of drinking water during 2 - 4 days.

Medicated drinking water should be used within 24 hours.

WITHDRAWAL PERIOD:

None.

STORAGE:

Store in a dry, dark place between 2 °C and 25 °C.

Store in closed packing.

Keep medicine away from children.

PACKING:

100 ml PE-Pharma bottle (144 bottles per box).

1 litre PE-Pharma bottle (12 bottles per box).