



POWERVIT WSP

COMPOSITION:

Contains per g:

Vitamin A	25,000	IU	Nicotinic acid	75	mg
Vitamin D	3,000	IU	Vitamin C	50	mg
Vitamin E	50	mg	DL-Methionine	10	mg
Vitamin K3	10	mg	L-Lysine	25	mg
Vitamin B1	10	mg	Manganese sulphate	10	mg
Vitamin B2	10	mg	Zinc sulphate	10	mg
Vitamin B6	10	mg	Iron sulphate	5	mg
Vitamin B12	25	µg	Copper sulphate	1.5	mg
Folic acid	0.2	mg	Cobalt sulphate	0.1	mg
Ca d-pantothenate	25	mg	Potassium iodide	0.1	mg

DESCRIPTION:

Powervit wsp is a scientifically concentrated, stabilised powder with multivitamins like the fat-soluble A, D, E and K and water-soluble vitamins of the B and C group; the essential amino acids methionine and lysine; and various essential minerals and trace elements like iron, copper, zinc, manganese, cobalt and iodide.

INDICATIONS:

Powervit wsp is indicated for stressful periods during the first days of life, vaccination, diseases, excessive changes in temperature, transfer to a new hen house, culling, transport etc. Post-treatment use after a Coccidiosis treatment; worm, bacterial and virus infections. Convalescence: during off-feed period and during deficiency or nutritional diseases.

Powervit wsp has also the following benefits for laying birds:

- Higher egg production peak level.
- Maintenance of high production level throughout the laying period.

SIDE EFFECTS:

None.

DOSAGE AND ADMINISTRATION:

For oral administration via feed or drinking water.

Poultry : 100 g per 400 litres of drinking water daily, during 1 - 3 days. The dosage may vary from 30 - 200 g per 400 litres according to individual requirements.

Mixed through feed: 500 - 1,000 g per 1,000 kg of feed.

Calves : 0.5 - 1 g per animal daily, during 7 days.

Pigs : 100 g per 200 - 400 litres of drinking water daily, during 3 - 5 days.

Mixed with feed: 500 - 1000 g per 1,000 kg of feed. Dosing depends on the individual requirements.

Sheep, goats : 0.5 g per animal daily, during 7 days.

Mixed with feed, the product should be used immediately. Medicated drinking water should be used within 24 hours.

WITHDRAWAL PERIOD:

None.

STORAGE:

Store in a dry, dark place between 2 °C and 25 °C.

Store in closed packing.

Keep medicine away from children.

PACKING:

Sachet of 100 g (120 sachets per box).

PE-Pharma jar of 500 / 1,000 g (12 jars per box).