



## FASCIONIX 25% INJ.

### COMPOSITION:

Contains per ml:

Nitroxynil 250 mg

### DESCRIPTION:

Nitroxynil is very effective for treatment of fascioliasis (mature and immature *Fasciola hepatica*) in cattle, sheep and goats. Nitroxynil is also very effective against *Haemonchus contortus* (adult and larval infestations with sheep and goats), *Bunostomum phlebotomum*, *Haemonchus plucei* and *Oesophagostomum radiatum* in cattle.

### INDICATIONS:

Fascionix 25% inj. is effective against:

- Liver fluke infestations caused by *Fasciola hepatica* and *Fasciola gigantica*. Gastrointestinal parasitism caused by *Haemonchus*, *Oesophagostomum* and *Bunostomum* in cattle, sheep and goats.
- *Oestrus ovis* in sheep and camels.
- Hookworms (*Ancylostoma* and *Uncinaria*) in dogs.

### CONTRA-INDICATIONS:

Do not use in animals in lactation if milk is intended for human consumption.

### SIDE EFFECTS:

None.

### WARNING:

Avoid contact with skin and mucosa.

### DOSAGE AND ADMINISTRATION:

For subcutaneous or oral administration.

- Cattle, sheep, goats, camels : 1 ml per 25 kg bodyweight by subcutaneous injection.  
This dosage may be increased to 1.25 ml per 25 kg bodyweight in case of acute Fascioliasis (immature flukes).  
: 5 ml per 10 litres of drinking water by oral administration.
- Dogs : 0.40 ml per 10 kg bodyweight by subcutaneous injection.  
: 0.60 ml per 10 kg bodyweight by oral administration.

No more than 20 ml should be injected into a single site. Repeat treatment only on advice of your veterinarian.

### WITHDRAWAL PERIOD:

- Meat: Cattle : 60 days  
Sheep, goats : 49 days

### STORAGE:

Store in a dry, dark place between 15 °C and 25 °C.

Keep medicine away from children.

### PACKING:

50 ml multidose vial packed in individual box (105 vials per box).

250 ml multidose vial packed in individual box (15 vials per box).