



FASCIONIX 34% INJ.

COMPOSITION:

Contains per ml:

Nitroxynil 340 mg

DESCRIPTION:

Nitroxynil is very effective for treatment of fascioliasis (mature and immature *Fasciola hepatica*) in cattle, sheep and goats. Nitroxynil is also very effective against *Haemonchus contortus* (adult and larval infestations with sheep and goats), *Bunostomum phlebotomum*, *Haemonchus plucei* and *Oesophagostomum radiatum* in cattle.

INDICATIONS:

Fascionix 34% Inj. is effective against:

- Liver fluke infestations caused by *Fasciola hepatica* and *Fasciola gigantica*. Gastrointestinal parasitism caused by *Haemonchus*, *Oesophagostomum* and *Bunostomum* in cattle, sheep and goats.
- *Oestrus ovis* in sheep and camels.
- Hookworms (*Ancylostoma* and *Uncinaria*) in dogs.

CONTRA-INDICATIONS:

Do not use in animals in lactation if milk is intended for human consumption.

SIDE EFFECTS:

None.

WARNING:

Avoid contact with skin and mucosa.

DOSAGE AND ADMINISTRATION:

For subcutaneous or oral administration.

- Cattle, sheep, goats, camels : 1 ml per 35 kg bodyweight by subcutaneous injection.
This dosage may be increased to 1.25 ml per 35 kg bodyweight in case of acute Fascioliasis (immature flukes).
: 4 ml per 10 litres of drinking water by oral administration.
- Dogs : 0.30 ml per 10 kg bodyweight by subcutaneous injection.
: 0.45 ml per 10 kg bodyweight by oral administration.

No more than 20 ml should be injected into a single site. Repeat treatment only on advice of your veterinarian.

WITHDRAWAL PERIOD:

- Meat: Cattle : 60 days
Sheep, goats : 49 days

STORAGE:

Store in a dry, dark place between 15 °C and 25 °C.

Keep medicine away from children.

PACKING:

50 ml multidose vial packed in individual box. (105 vials per box).

100 ml multidose vial packed in individual box. (48 vials per box).